

PRACTICE QUESTION PAPER - IV

CLASS XII - MATHEMATICS (041)

Time Allowed: 3 Hours

Maximum Marks: 80

General Instructions:

1. This Question Paper contains **38** questions. All questions are compulsory.
 2. The question paper is divided into FIVE Sections – A, B, C, D and E.
 3. Section **A** comprises of **20** questions of **1** mark each. (18 MCQs + 2 Assertion-Reasoning)
 4. Section **B** comprises of **5** questions of **2** marks each.
 5. Section **C** comprises of **6** questions of **3** marks each.
 6. Section **D** comprises of **4** questions of **5** marks each.
 7. Section **E** comprises of **3** Case Study Based Questions of **4** marks each.
 8. There is no overall choice in the question paper. However, an internal choice has been provided in **2** questions in Section B, **3** questions in Section C, **2** questions in Section D and **2** questions in Section E (in the sub-parts).
 9. Use of calculators is **not** permitted.
-

SECTION A (20 Marks)

This section comprises 20 questions of 1 mark each. Questions 1-18 are Multiple Choice Questions (MCQs) and questions 19-20 are Assertion-Reason based questions.

Multiple Choice Questions (MCQs)

1. The relation $R = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 3)\}$ on set $A = \{1, 2, 3\}$ is:
 - (a) Only Reflexive
 - (b) Only Transitive
 - (c) Reflexive and Transitive
 - (d) An Equivalence relation
2. If $f : \mathbb{R} \rightarrow \mathbb{R}$ is defined by $f(x) = |x|$, then the range of f is:
 - (a) \mathbb{R}
 - (b) $[0, \infty)$
 - (c) $(-\infty, 0)$
 - (d) $(0, \infty)$
3. The value of $\cos(\sec^{-1} x + \csc^{-1} x)$ for $|x| \geq 1$ is:
 - (a) 1
 - (b) -1
 - (c) 0
 - (d) $\frac{\pi}{2}$
4. If $\tan^{-1} x + \tan^{-1} y = \frac{\pi}{4}$ and $xy < 1$, then $x + y + xy$ is equal to:
 - (a) 1
 - (b) 0
 - (c) -1
 - (d) $\frac{1}{2}$

5. The set $A = \{a, b, c\}$ and $B = \{1, 2\}$. The number of onto functions from A to B is:

- (a) 6
- (b) 7
- (c) 2
- (d) 8

6. If the matrix $A = \begin{bmatrix} 0 & a \\ -a & 0 \end{bmatrix}$, then A^3 is equal to:

- (a) $\begin{bmatrix} 0 & a^3 \\ -a^3 & 0 \end{bmatrix}$
- (b) $\begin{bmatrix} 0 & -a^3 \\ a^3 & 0 \end{bmatrix}$
- (c) $\begin{bmatrix} a^3 & 0 \\ 0 & a^3 \end{bmatrix}$
- (d) $\begin{bmatrix} -a^3 & 0 \\ 0 & -a^3 \end{bmatrix}$

7. If A is an invertible matrix and $A^{-1} = \begin{bmatrix} 3 & -1 \\ -4 & 1 \end{bmatrix}$, then the matrix A is:

- (a) $\begin{bmatrix} 1 & 1 \\ 4 & 3 \end{bmatrix}$
- (b) $\begin{bmatrix} -1 & -1 \\ -4 & -3 \end{bmatrix}$
- (c) $\begin{bmatrix} 1 & -1 \\ -4 & 3 \end{bmatrix}$
- (d) $\begin{bmatrix} 1 & 4 \\ 1 & 3 \end{bmatrix}$

8. If A is a square matrix of order n , then $|\text{adj}(\text{adj}(A))|$ is:

- (a) $|A|^{n-1}$
- (b) $|A|^{n(n-1)}$
- (c) $|A|^{(n-1)^2}$
- (d) $|A|^{n^2}$

9. If A is a 3×3 matrix and $\text{adj}(A) = \begin{bmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 5 \end{bmatrix}$, then $|A|$ is:

- (a) 5
- (b) 25
- (c) 125
- (d) -5

10. Let A be a non-zero column matrix. The rank of AA^T is:

- (a) 0
- (b) 1
- (c) 2
- (d) 3

11. If $y = \log(x + \sqrt{x^2 + a^2})$, then $\frac{dy}{dx}$ is:
- $\frac{1}{\sqrt{x^2 + a^2}}$
 - $\frac{1}{x + \sqrt{x^2 + a^2}}$
 - $\frac{x}{\sqrt{x^2 + a^2}}$
 - $\sqrt{x^2 + a^2}$
12. The particular solution of the differential equation $\frac{dy}{dx} = 2x$ when $y(0) = 0$ is:
- $y = x^2$
 - $y = x^2 + 1$
 - $y = 2x^2$
 - $y = 2x^2 + 1$
13. The rate of change of area of a circle with respect to its diameter is:
- $2\pi r$
 - πr
 - π
 - 2π
14. The maximum value of the function $f(x) = 3 - 2\sin x$ is:
- 5
 - 3
 - 1
 - 1
15. The value of $\int \frac{1}{\sin^2 x \cos^2 x} dx$ is:
- $\tan x + \cot x + C$
 - $\tan x - \cot x + C$
 - $\sec x - \csc x + C$
 - $-\tan x + \cot x + C$
16. The degree of the differential equation $\left[1 + \left(\frac{dy}{dx}\right)^2\right]^{3/2} = \frac{d^2y}{dx^2}$ is:
- 1
 - 2
 - 3
 - Not defined
17. The coordinates of the foot of the perpendicular from the origin on the plane $2x - 3y + 4z - 6 = 0$ are:
- $\left(\frac{12}{29}, -\frac{18}{29}, \frac{24}{29}\right)$
 - $(2, -3, 4)$
 - $(12, -18, 24)$
 - $\left(\frac{12}{13}, -\frac{18}{13}, \frac{24}{13}\right)$
18. If the scalar projection of vector \vec{a} on vector \vec{b} is $|\vec{b}|$, then the angle between \vec{a} and \vec{b} is:
- 0

- (b) $\frac{\pi}{6}$
- (c) $\frac{\pi}{4}$
- (d) $\frac{\pi}{3}$

Assertion-Reasoning Based Questions

Questions 19 and 20 are Assertion-Reasoning based questions. In these questions, a statement of Assertion (A) is followed by a statement of Reason (R). Choose the correct answer from the following options:

- (a) Both A and R are true and R is the correct explanation of A.
 - (b) Both A and R are true but R is not the correct explanation of A.
 - (c) A is true but R is false.
 - (d) A is false but R is true.
19. **Assertion (A):** The function $f(x) = x^3 - 6x^2 + 15x - 8$ is strictly increasing on \mathbb{R} . **Reason (R):** A differentiable function $f(x)$ is strictly increasing if $f'(x) > 0$ for all x .
20. **Assertion (A):** If A and B are two matrices such that AB and BA are defined, then $AB = BA$. **Reason (R):** Matrix multiplication is generally commutative.
-

SECTION B (10 Marks)

This section comprises 5 questions of 2 marks each.

- 21. Find the value of $\frac{dy}{dx}$ at $t = \frac{\pi}{2}$ if $x = a(\cos t + t \sin t)$ and $y = a(\sin t - t \cos t)$.
- 22. Find the area of the triangle whose vertices are $A(1, 1, 1)$, $B(1, 2, 3)$, and $C(2, 3, 1)$.

OR

Show that the lines $\frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$ and $\frac{x-3}{-4} = \frac{y-2}{-3} = \frac{z-1}{-2}$ intersect.

- 23. Find $\int \sin^{-1}(\cos x) dx$.

OR

Find $\int \frac{dx}{\sqrt{7-6x-x^2}}$.

- 24. Prove that the function $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = \sin x$ is neither one-one nor onto.
 - 25. A die is tossed twice. Let E be the event "first toss shows a 5" and F be the event "sum of outcomes is greater than 9". Find $P(E|F)$.
-

SECTION C (18 Marks)

This section comprises 6 questions of 3 marks each.

- 26. Simplify $\tan^{-1} \left(\frac{\sqrt{1+x^2}-1}{x} \right)$, where $x \neq 0$.
- 27. Find the maximum value of $4x + \frac{16}{x}$ for $x > 0$.

OR

Find the derivative of $\sin x$ w.r.t. x^2 .

- 28. Find the general solution of the differential equation $\cos^2 x \frac{dy}{dx} + y = \tan x$.

OR

Show that $y = Ax + B/x$ is a solution of the differential equation $x^2 \frac{d^2y}{dx^2} + x \frac{dy}{dx} - y = 0$.

- 29. Find the volume of the parallelepiped whose coterminal edges are represented by the vectors $\vec{a} = 2\hat{i} - 3\hat{j} + 4\hat{k}$, $\vec{b} = \hat{i} + 2\hat{j} - \hat{k}$ and $\vec{c} = 3\hat{i} - \hat{j} + 2\hat{k}$.

OR

Find the equation of the plane through the intersection of the planes $3x - y + 2z - 4 = 0$ and $x + y + z - 2 = 0$ and passing through the point $(2, 2, 1)$.

30. Using properties of determinants, show that $\begin{vmatrix} 1 & x & x^2 \\ x^2 & 1 & x \\ x & x^2 & 1 \end{vmatrix} = (1 - x^3)^2$.

31. Minimize the objective function $Z = 5x + 10y$ subject to $x + 2y \leq 120$, $x + y \geq 60$, $x - 2y \geq 0$, and $x, y \geq 0$. (Identify the corner points only).

SECTION D (20 Marks)

This section comprises 4 questions of 5 marks each.

33. Find the area of the region bounded by the curves $y^2 = 4x$ and $x^2 = 4y$.

OR

Find the area bounded by the curve $y = \sqrt{x}$ and the line $y = x$.

34. If $A = \begin{bmatrix} 1 & -1 & 1 \\ 1 & -2 & -2 \\ 2 & 1 & 3 \end{bmatrix}$, find A^{-1} . Hence, solve the system of linear equations:

$$\begin{aligned} x - y + z &= 4 \\ x - 2y - 2z &= 9 \\ 2x + y + 3z &= 1 \end{aligned}$$

35. A window is in the form of a rectangle surmounted by a semicircle. The total perimeter of the window is 10 m. Find the dimensions of the window to admit maximum light through the whole opening.

OR

Evaluate $\int \frac{x^2+1}{(x^2+4)(x^2+2)} dx$.

36. Find the shortest distance between the lines $\vec{r} = (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda(\hat{i} - 3\hat{j} + 2\hat{k})$ and $\vec{r} = (4\hat{i} + 5\hat{j} + 6\hat{k}) + \mu(2\hat{i} + 3\hat{j} + \hat{k})$.

SECTION E (12 Marks)

This section comprises 3 case study based questions of 4 marks each.

37. Case Study 1: Drone Trajectory and Collinearity

A surveillance drone is flying in a path defined by a line segment AB connecting two points $A(1, 2, -1)$ and $B(4, 6, 8)$. A sensor is located at point $C(10, 14, 26)$.

Based on the given information, answer the following questions:

- (a) Find the vector \vec{AB} . (1 Mark)
- (b) Determine if the points A, B , and C are collinear. (1 Mark)
- (c) Find the direction cosines of the line segment BC . (2 Marks)

OR

- (d) Find the vector component of \vec{AC} perpendicular to \vec{AB} . (2 Marks)

38. Case Study 2: Life Insurance and Expected Value

A life insurance company offers a policy where, for an annual premium of Rs.500, a person gets Rs.10,000 if they die within the year. The probability of a person dying in the age group covered by the policy is 0.005.

Let X be the random variable representing the gain (or loss) for the insurance company.

Based on the given information, answer the following questions:

- (a) What is the value of X if the person survives the year? (1 Mark)
- (b) Prepare the probability distribution of X . (1 Mark)
- (c) Calculate the expected gain (or loss) $E(X)$ for the insurance company on one policy. (2 Marks)

OR

- (d) Calculate the variance of X (You may use $E(X^2) = 50000$). (2 Marks)

39. Case Study 3: Rate of Change and Volume

A storage tank is in the shape of a cube. The side length of the cube is increasing at a constant rate of 5 cm/s due to expansion. The current side length is 10 cm.

Based on the given information, answer the following questions:

- (a) Write the formula for the volume V of the cube in terms of its side a . (1 Mark)
- (b) Find the rate at which the volume of the cube is increasing ($\frac{dV}{dt}$) when the side length is 10 cm. (3 Marks)

OR

- (c) If the rate of change of volume were proportional to the square of the side, i.e., $\frac{dV}{dt} = ka^2$, find the value of the constant k . (3 Marks)
-